

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 November 2003 (06.11.2003)

PCT

(10) International Publication Number
WO 03/090737 A2

(51) International Patent Classification⁷: **A61K 31/196,**
C07C 229/42

WC1E 6BN (GB). **GONZALEZ, Isabel** [ES/GB]; 55 St
Barnabas Road, Cambridge, CB1 2BX (GB).

(21) International Application Number: PCT/EP03/04152

(74) Agent: **GRUBB, Philip**; Novartis AG, Corporate Intellect-
tual Property, CH-4002 Basel (CH).

(22) International Filing Date: 22 April 2003 (22.04.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0209257.5 23 April 2002 (23.04.2002) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LT, LU, LV, MA, MD, MK, MN, MX, NI, NO, NZ, OM,
PH, PL, PT, RO, RU, SC, SE, SG, SK, TJ, TM, TN, TR,
TT, UA, US, UZ, VC, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except AT, US*): **NO-
VARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel
(CH).

(84) Designated States (*regional*): Eurasian patent (AM, AZ,
BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE,
BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU,
IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR).

(71) Applicant (*for AT only*): **NOVARTIS PHARMA GMBH**
[AT/AT]; Brunner Strasse 59, A-1230 Viena (AT).

Published:


— without international search report and to be republished
upon receipt of that report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **FUJIMOTO,**
Roger, Aki [US/US]; 21 Molly Stark Drive, Morristown,
NJ 07960 (US). **URBAN, Laszlo** [GB/GB]; The Novartis
Institute for Medical Sciences, 5 Gower Place, London,

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: METHODS AND COMPOSITIONS FOR TREATMENT OF CANCER PAIN



(57) Abstract: The invention provides a method of treating cancer pain, e.g. bone cancer pain, in a subject in need of such treatment which comprises administering to the subject an effective amount of a COX-2 inhibitor; advantageously a compound of formula (I); wherein R, R₁, R₂, R₃, R₄, and R₅ are as defined; a pharmaceutically acceptable salts thereof; or a pharmaceutically acceptable prodrug ester thereof.

METHODS AND COMPOSITIONS FOR TREATMENT OF CANCER PAIN

This invention relates to selective cyclooxygenase-2 inhibitors (COX-2 Inhibitors), in particular to the use of COX-2 inhibitors in the treatment of cancer pain, especially bone cancer pain.

COX-2 inhibitors and their use as non-steroidal anti-inflammatory drugs (NSAIDs) for the treatment of inflammatory diseases and pain are well known in the art. Further it has been proposed (WO 99/11605) that COX-2 inhibitors may be useful for treatment of neoplasia particularly neoplasias that produce prostaglandins or express cyclooxygenase, including both benign and cancerous tumors, growths and polyps.


It has now been found that certain COX-2 inhibitors, in particular 5-alkyl substituted 2-arylamino-phenylacetic acid derivative COX-2 inhibitors, have desirable properties for use in the treatment of Cancer pain, in particular bone cancer pain.

Accordingly the invention provides a method of treating cancer pain in a subject in need of such treatment, which comprises administering to the subject an effective amount of a COX-2 inhibitor.

Suitable COX-2 inhibitors for use in the invention may include the following compounds or derivatives thereof or a pharmaceutically acceptable salt thereof, or any hydrate thereof: rofecoxib, etoricoxib, celecoxib, valdecoxib, parecoxib, or a 5-alkyl-2-arylamino-phenylacetic acid derivative COX-2 inhibitor.

In preferred embodiments the invention provides a method of treating cancer pain in a subject in need of such treatment which comprises administering to the subject an effective amount of a COX-2 inhibitor of formula I

- 2 -



wherein R is methyl or ethyl;

R₁ is chloro or fluoro;

R₂ is hydrogen or fluoro;

R₃ is hydrogen, fluoro, chloro, methyl, ethyl, methoxy, ethoxy or hydroxy;

R₄ is hydrogen or fluoro; and

R₅ is chloro, fluoro, trifluoromethyl or methyl;

a pharmaceutically acceptable salt thereof; or

a pharmaceutically acceptable prodrug ester thereof.

Further the invention provides the use of a compound of formula I (or pharmaceutically acceptable salt or prodrug ester thereof) as defined above for the preparation of a medicament, for use in the treatment of cancer pain.

In a further aspect the invention provides use of a Cox-2 inhibitor, advantageously a compound of formula I (or pharmaceutically acceptable salt or prodrug ester thereof) as defined above for the treatment of cancer pain.

In a yet further aspect the invention provides cancer pain treatment agent comprising a Cox-2 inhibitor, advantageously a compound of formula I (or pharmaceutically acceptable salt or prodrug ester thereof) as defined above as active ingredient.

In a still yet further aspect the invention provides a package comprising a Cox-2 inhibitor, advantageously a compound of formula I (or pharmaceutically acceptable salt or

prodrug ester thereof) as defined above together with instructions for use in the treatment of cancer pain.

The compounds of formula I may be used for treatment of cancer pain in general. In a particularly preferred embodiment the compounds of formula I are used for the treatment of bone cancer pain including pain associated with primary bone cancer such osteosarcoma, and also pain associated with bone metastases of primary cancers such as breast, colon, lung, prostate and other cancers, e.g. multiple myeloma.

In the present description the terms "treatment" or "treat" refer to both prophylactic or preventative treatment as well as curative or disease modifying treatment, including treatment of patients at risk of contracting the disease or suspected to have contracted the disease as well as patients who are ill or have been diagnosed as suffering from a disease or medical condition.

Particularly preferred compounds of formula I are those wherein R is methyl or ethyl; R₁ is chloro or fluoro; R₂ is hydrogen; R₃ is hydrogen, fluoro, chloro, methyl or hydroxy; R₄ is hydrogen; and R₅ is chloro, fluoro or methyl; pharmaceutically acceptable salts thereof; and pharmaceutically acceptable esters thereof.

A particularly preferred embodiment relates to the compounds of formula I wherein R is methyl or ethyl; R₁ is fluoro; R₂ is hydrogen; R₃ is hydrogen, fluoro or hydroxy; R₄ is hydrogen; and R₅ is chloro; pharmaceutically acceptable salts thereof; and pharmaceutically acceptable prodrug esters thereof.

Another particularly preferred embodiment of the invention relates to compounds of formula I wherein R is ethyl or methyl; R₁ is fluoro; R₂ is hydrogen or fluoro; R₃ is hydrogen, fluoro, ethoxy or hydroxy; R₄ is hydrogen or fluoro; and R₅ is chloro, fluoro or methyl; pharmaceutically acceptable salts thereof; and pharmaceutically acceptable prodrug esters thereof.

Further are said compounds wherein R is methyl or ethyl; R₁ is fluoro; R₂-R₄ are hydrogen or fluoro; and R₅ is chloro or fluoro; pharmaceutically acceptable salts thereof; and pharmaceutically acceptable prodrug esters thereof.

A further embodiment of the invention relates to the compounds of formula I wherein R is methyl or ethyl; R₁ is fluoro; R₂ is fluoro; R₃ is hydrogen, ethoxy or hydroxy; R₄ is fluoro; and R₅ is fluoro; pharmaceutically acceptable salts thereof; and pharmaceutically acceptable prodrug esters thereof.

Another embodiment of the invention relates to the compounds of formula I wherein R is methyl; R₁ is fluoro; R₂ is hydrogen; R₃ is hydrogen or fluoro; R₄ is hydrogen; and R₅ is chloro; pharmaceutically acceptable salts thereof; and pharmaceutically acceptable prodrug esters thereof.

Particularly preferred embodiments of the invention relate to compounds of formula I

(a) wherein R is methyl; R₁ is fluoro; R₂ is hydrogen; R₃ is hydrogen; R₄ is hydrogen; and R₅ is chloro; pharmaceutically acceptable salts thereof; and pharmaceutically acceptable prodrug esters thereof;

(b) wherein R is methyl; R₁ is fluoro; R₂ is hydrogen; R₃ is fluoro; R₄ is hydrogen; and R₅ is chloro; pharmaceutically acceptable salts thereof; and pharmaceutically acceptable prodrug esters thereof;


(c) wherein R is ethyl; R₁ is fluoro; R₂ is fluoro; R₃ is hydrogen; R₄ is fluoro; and R₅ is fluoro; pharmaceutically acceptable salts thereof; and pharmaceutically acceptable prodrug esters thereof; and

(d) wherein R is ethyl; R₁ is chloro; R₂ is hydrogen; R₃ is chloro; R₄ is hydrogen; and R₅ is methyl; pharmaceutically acceptable salts thereof; and pharmaceutically acceptable prodrug esters thereof.

Most preferably 5-methyl-2-(2'-chloro-6'-fluoroanilino)-phenylacetic acid or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable prodrug thereof is used as the COX-2 inhibitor of the invention.

Pharmacologically acceptable salts of the compounds of formula I are preferably salts with bases, conveniently metal salts derived from groups Ia, Ib, IIa and IIb of the Periodic Table of the Elements, including alkali metal salts, e.g. potassium and especially sodium salts, or alkaline earth metal salts, preferably calcium or magnesium salts, and also ammonium salts with ammonia or organic amines.

Pharmaceutically acceptable prodrug esters of the compounds of formula I are ester derivatives which are convertible by solvolysis or under physiological conditions to the free carboxylic acids of formula I. Such esters are e.g. lower alkyl esters (such as the methyl or ethyl ester), carboxy-lower alkyl esters such as the carboxymethyl ester, nitrooxy-lower alkyl esters (such as the 4-nitrooxybutyl ester), and the like. Preferred prodrugs are the compounds of formula Ia



wherein R and R₁.R₅ have meaning as defined hereinabove for compounds of formula I; and pharmaceutically acceptable salts thereof.

Compounds of formula I and Ia and their synthesis are described in published international patent applications Nos. WO 99/11605 and WO 01/23346, the teachings of which are incorporated herein by reference.

It has been further discovered in accordance with the present invention that compounds of formula I (and esters and prodrugs thereof), in particular the compound 5-methyl-2-(2'-chloro-6'-fluoroanilino)-phenylacetic acid or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable prodrug thereof, inhibit the radiologically observed structural changes in an animal model of bone cancer pain.

Thus in a yet further embodiment the invention includes a method for the inhibition of bone loss, advantageously in cancer, which comprises administering an effective amount of a COX-2 inhibitor of formula I (or an ester or prodrug thereof) as defined above to the subject in need of such treatment.

The Agents of the Invention, i.e. the COX-2 inhibitors of formula I and pharmaceutically acceptable salts and esters thereof , are preferably used in the form of pharmaceutical preparations that contain the relevant therapeutically effective amount of active ingredient optionally together with or in admixture with inorganic or organic, solid or liquid, pharmaceutically acceptable carriers which are suitable for administration.

The COX-2 pharmaceutical compositions may be, for example, compositions for enteral, such as oral, rectal, aerosol inhalation or nasal administration; compositions for parenteral, such as intravenous or subcutaneous administration; compositions for transdermal administration (e.g. passive or iontophoretic); or compositions for topical administration.

Preferably, the COX-2 pharmaceutical compositions are adapted to oral or parenteral (especially oral) administration. Intravenous and oral, first and foremost oral, administration is considered to be of particular importance. Preferably the COX-2 inhibitor active ingredient is in oral form.

The particular mode of administration and the dosage may be selected by the attending physician taking into account the particulars of the patient, especially age, weight, life style, activity level, etc .

The dosage of the Agents of the Invention may depend on various factors, such as effectiveness and duration of action of the active ingredient, mode of administration, warm-blooded species, and/or sex, age, weight and individual condition of the warm-blooded animal.

More particularly, the pharmaceutical compositions comprise an effective cyclooxygenase-2 inhibiting amount of COX-2 inhibitor or compound of formula I which is substantially free of cyclooxygenase-1 inhibiting activity and of side effects attributed thereto, when used therapeutically.

The compounds of formula I (and salts and esters thereof) are useful in the manufacture of pharmaceutical compositions comprising an effective amount thereof in conjunction or admixture with excipients or carriers suitable for either enteral or parenteral application. Preferred are tablets and gelatin capsules comprising the active ingredient together with a) diluents, e.g. lactose, dextrose, sucrose, mannitol, sorbitol, cellulose and/or glycine; b) lubricants, e.g. silica, talcum, stearic acid, its magnesium or calcium salt and/or polyethyleneglycol; for tablets also c) binders e.g. magnesium aluminum silicate, starch paste, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose and or polyvinylpyrrolidone; if desired d) disintegrants, e.g. starches, agar, alginic acid or its sodium salt, or effervescent mixtures; and/or e) absorbents, colorants, flavors and sweeteners. Injectable compositions are preferably aqueous isotonic solutions or suspensions, and suppositories are advantageously prepared from fatty emulsions or suspensions. Said compositions may be sterilized and/or contain adjuvants, such as preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for regulating the osmotic pressure and/or buffers. In addition, they may also contain other therapeutically valuable substances. Said compositions are prepared according to conventional mixing, granulating or coating

methods, respectively, and contain about 0.1 to 80%, preferably about 1 to 60% or more, of the active ingredient.

Tablets may be either film coated or enteric coated according to methods known in the art.

Suitable formulations for transdermal application include an effective amount of a compound of the invention with carrier. Advantageous carriers include absorbable pharmacologically acceptable solvents to assist passage through the skin of the host. For example, transdermal devices are in the form of a bandage comprising a backing member, a reservoir containing the compound optionally with carriers, optionally a rate controlling barrier to deliver the compound of the skin of the host at a controlled and predetermined rate over a prolonged period of time, and means to secure the device to the skin.

Suitable formulations for topical application, e.g. to the skin and eyes, include aqueous solutions, suspensions, ointments, creams, gels or sprayable formulations, for example, for delivery by aerosol or the like. Such topical delivery systems will in particular be appropriate for dermal application, e.g. for the treatment of skin cancer, for example, for prophylactic use in creams, lotions sprays and the like

The dosage of COX-2 inhibitor administered is dependent on the species of warm-blooded animal (mammal), the body weight, age and individual condition, and on the form of administration. A unit dosage for oral administration to a mammal of about 50 to 70 kg may contain between about 5 and 1500 mg, e.g. from 100-1000 mg, preferably 200-800 mg of the active ingredient.

COX-2 inhibitor formulations in single dose unit form contain preferably from about 1% to about 90%, and formulations not in single dose unit form contain preferably from about 0.1% to about 20%, of the active ingredient. Single dose unit forms such as capsules, tablets or dragées contain e.g. from about 1mg to about 1000mg of the active ingredient.

COX-2 inhibitor pharmaceutical preparations for enteral and parenteral administration are, for example, those in dosage unit forms, such as dragées, tablets or capsules and also ampoules. They are prepared in a manner known *per se*, for example by means of conventional mixing, granulating, confectioning, dissolving or lyophilising processes. For example, pharmaceutical preparations for oral administration can be obtained by combining the active ingredient with solid carriers, where appropriate granulating a resulting mixture, and processing the mixture or granulate, if desired or necessary after the addition of suitable adjuncts, into tablets or dragée cores.

Parenteral formulations are especially injectable fluids that are effective in various manners, such as intravenously, intramuscularly, intraperitoneally, intranasally, intradermally or subcutaneously. Such fluids are preferably isotonic aqueous solutions or suspensions which can be prepared before use, for example from lyophilised preparations which contain the active ingredient alone or together with a pharmaceutically acceptable carrier. The pharmaceutical preparations may be sterilised and/or contain adjuncts, for example preservatives, stabilisers, wetting agents and/or emulsifiers, solubilisers, salts for regulating the osmotic pressure and/or buffers.

The following examples are intended to illustrate the invention and are not to be construed as being limitations thereon.

EXAMPLES**A. Formulation Examples****Example 1**

Table 1

Ingredient	Amount per 200 mg tablet batch (kg)
Core	
Granulation	
5-methyl-2-(2'-chloro-6'-fluoroanilino)phenylacetic acid drug substance	50**
Microcrystalline cellulose, NF (PH 101)	12.85
Lactose monohydrate, NF	11.65
Croscarmellose sodium, NF	1
Povidone, USP	4
Titanium dioxide, USP	2
Water, purified ***, USP	20.375
Extra-granular Phase	
Microcrystalline cellulose, NF (PH 102)	13
Croscarmellose sodium, NF	3
Titanium dioxide, USP	2
Magnesium stearate, NF	0.5
Coating	
Opadry white	2.801 ****
Opadry yellow	2.0 ****
Opadry red	0.4 ****
Opadry black	0.0504 ****
Water, purified ***, USP	29.758 ****

** The weight of drug substance is taken with reference to the dried substance (100 per cent) on the basis of the assay value (factorization). The difference in weight is adjusted by the amount of microcrystalline cellulose used.

*** Removed during processing.

**** Includes a 50 % excess for loss during the coating process.

Table 1, above, sets out the formula for a batch of approximately 250,000 immediate release film-coated tablets of 5-methyl-2-(2'-chloro-6'-fluoroanilino)-phenylacetic acid. To make the tablets, titanium dioxide is dispersed in water, followed by the addition of povidone and mixing for 20 minutes to make a povidone/titanium dioxide suspension. The drug substance, lactose, microcrystalline cellulose, and croscarmellose are mixed in a high shear mixer (e.g., a Collette Gral) for 5 minutes to form a drug mixture. The drug mixture is granulated in the high shear mixer with the povidone/titanium dioxide suspension. The suspension is pumped at a rate of 3 kg/min into the drug mixture. The resulting mixture is mixed an additional 90 seconds after all the suspension is added. The wet granulation is dried in a fluid bed dryer, using an inlet air temperature of 50 °C. The residual water target is 3.5 % (with a permissible range of 2.5 – 4.5 %). The dried granulation is passed through a screen using a mill (oscillator) and a 30 mesh screen. The previous steps are repeated to make a second granulation.

The extra-granular phase titanium dioxide is passed through a 60 mesh hand screen. The dry granulations are mixed with the extra-granular phase microcrystalline cellulose, croscarmellose sodium and titanium dioxide in a twin shell mixer for 300 revolutions to form a penultimate mixture. Magnesium stearate is passed through a 60 mesh hand screen and is mixed with the penultimate mixture in a twin shell mixer for 50 revolutions to form a tableting mixture. The tableting mixture is pressed into tablets using a tablet press and oval punches.

The coating powders (Opadry) are mixed with purified water to make a 15 % w/w coating suspension. The tablets are film coated with the coating suspension in a coating pan using 60 °C to 75 °C inlet air temperature.

Table 2 sets out the contents of a 200 mg 5-methyl-2-(2'-chloro-6'-fluoroanilino)phenylacetic acid film-coated tablet.

Table 2

Ingredient	Theoretical amount [mg]	Function
Core		
5-methyl-2-(2'-chloro-6'-fluoroanilino)phenylacetic acid drug substance	200	Active substance
Microcrystalline cellulose (PH 101)	51.4	Filler
Lactose	46.6	Filler
Povidone	16	Binder
Titanium dioxide	8	Color
Croscarmellose sodium	4	Disintegrant
Water, purified *	Q.S.	Granulating liquid
Extragranular phase		
Microcrystalline cellulose (PH 102)	52	Filler
Croscarmellose sodium	12	Disintegrant
Titanium dioxide	8	Color
Magnesium stearate	2	Lubricant
Core weight	400	
Coating		
Opadry white (00F18296)	7.4676	Color
Opadry yellow (00F12951)	5.3312	Color
Opadry red (00F15613)	1.0668	Color
Opadry black (00F17713)	0.1344	Color

- 13 -

Ingredient	Theoretical amount [mg]	Function
Water, purified *	Q.S.	Coating solvent
Total weight	414	

* removed during processing

In addition, the tablet formulations may contain 5-methyl-2-(2'-chloro-6'-fluoroanilino)benzyl alcohol and/or 5-methyl-2-(2'-chloro-6'-fluoroanilino)benzoic acid in an amount between about 0.01 and 2% by weight, more specifically between about 0.1 and 1.

Example 2

Wet granulated tablet composition

<u>Amount per tablet</u>	<u>Ingredient</u>
25 mg	COX-2 inhibitor
79.7 mg	Microcrystalline cellulose
79.7 mg	Lactose monohydrate
6 mg	Hydroxypropyl cellulose
8 mg	Croscarmellose sodium
0.6 mg	Iron oxide
1 mg	Magnesium stearate

Tablet dose strengths of between 5 and 125 mg can be accommodated by varying total weight, and the ratio of the first three ingredients. Generally it is preferable to maintain a 1:1 ratio for microcrystalline cellulose: lactose monohydrate.

Example 3

Directly compressed tablet composition

<u>Amount per tablet</u>	<u>Ingredient</u>
--------------------------	-------------------

- 14 -

25	mg	COX-2 inhibitor
106.9	mg	Microcrystalline cellulose
106.9	mg	Lactose anhydrate
7.5	mg	Croscarmellose sodium
3.7	mg	Magnesium stearate

Tablet dose strengths of between 5 and 125 mg can be accommodated by varying total tablet weight, and the ratio of the first three ingredients. Generally it is preferable to maintain a 1:1 ratio for microcrystalline cellulose:lactose monohydrate.

Example 4

Hard gelatine capsule composition

<u>Amount per capsule</u>	<u>Ingredient</u>
25 mg	COX-2 inhibitor
37 mg	Microcrystalline cellulose
37 mg	Lactose anhydrate
1 mg	Magnesium stearate
1 capsule	Hard gelatin capsule

Capsule dose strengths of between 1 and 50 mg can be accommodated by varying total fill weight, and the ratio of the first three ingredients. Generally it is preferable to maintain a 1:1 ratio for microcrystalline cellulose:lactose monohydrate.

Example 5

Oral solution

<u>Amount per 5mL</u>	<u>Ingredient</u>
50 mg	COX-2 inhibitor
to 5 mL	with Polyethylene oxide 400

Example 6

Oral suspension

Amount per 5mL dose Ingredient

101	mg	COX-2 inhibitor
150	mg	Polyvinylpyrrolidone

Oral suspensionAmount per 5mL dose Ingredient

2.5	mg	Poly oxyethylene sorbitan monolaurate
10	mg	Benzoic acid
to 5 mL with sorbitol solution (70%)		

Suspension dose strengths of between 1 and 50 mg/5 ml can be accommodated by varying the ratio of the first two ingredients.

Example 7Intravenous infusionAmount per 200 mL dose Ingredient

1	mg	COX-2 inhibitor
0.2	mg	Polyethylene oxide 400
1.8	mg	Sodium chloride
to 200 mL		Purified water

Example 8 The Effect of COX-2 Inhibitors in a Rat Model of Bone Cancer pain

Adult female rats are given intra-tibial injections of MRMT-1 rat mammary gland carcinoma cells (3 μ l, 10⁷ cells/ml). These animals gradually develop mechanical hyperalgesia, mechanical allodynia (skin sensitivity to non-noxious stimuli) and hind limb sparing, beginning on day 12-14 following cell injection. COX-2 inhibitors, 5-methyl-2-(2'-chloro-6'-

fluoroanilino)-phenylacetic acid (hereinafter referred to as COX189) (10 and 30 mg/kg p.o.) and valdecoxib (30 mg/kg p.o.) are administered as described below and the results obtained compared with vehicle treated controls.

Intra-tibial inoculation of MRMT-1 cells induces a time-dependent hypersensitivity in the ipsilateral limb, measured as a shift in body weight to the contralateral limb. The increase in weight-bearing difference between contralateral and ipsilateral paw (weight shift from the affected to the unaffected limb) is evident by day 14 post-inoculation and continues to increase until day 20.

The repeated administration of COX189 (10 and 30mg/kg, p.o.) from day 10 post-tumour cell inoculation significantly attenuates the weight-bearing difference on days 14, 17 and 20 measured 1 hour after drug administration. This effect is significant for COX189 on days 10, 14, 17 and 20 as well as for valdecoxib (30mg/kg, p.o.) on day 20. The doses chosen for COX189 and valdecoxib are comparable. Lower doses are not studied.

The effect is not dose-dependent, there is no difference between the effect of COX189 administered at doses of 10 and 30mg/kg. The treatments induces a progressive and long-lasting reduction in hypersensitivity, since the weight-bearing difference is also attenuated when measured 1h before the first daily treatment. This preventative effect reaches significant levels for all three treatments on days 14 and 17. There is no significant difference between the effects measured before and after COX189 administration on each day for any individual treatment.

There is pronounced mechanical allodynia in the ipsilateral paw (vehicle group) both to a punctuate stimulus (static allodynia) and to light stroking of the skin (dynamic allodynia) on day 20 post-inoculation. The repeated administration of COX189 (10 and 30mg/kg, p.o.) or valdecoxib (30mg/kg, p.o.) significantly reverses static allodynia on day 20, measured 90 minutes after the last administration. There is a tendency to reduce dynamic allodynia by all treatments, although this reduction only reaches statistical significance for COX189 at the highest dose (30mg/kg, p.o.).


Measurement of COX189 levels in serum of the same animals, taken 3 hours after administration on day 20, shows concentrations of mean \pm SEM: 19.0 \pm 1.7 μ M and 59.9 \pm 3.9 μ M for COX189, 10 and 30mg/kg, respectively

The levels of thromboxane B₂ in the samples from COX189-treated animals are 87.4±17.7%, 64.5±8.0% and 77.4±27.8% of the levels in the vehicle-control animals, for COX189 (10mg/kg), COX189 (30mg/kg) and valdecoxib (30mg/kg), respectively. These data indicate that COX-1 is not inhibited by these doses, implying that the benefits seen in the model are COX-2 as opposed to COX-1 derived.

In addition it is found in this animal model experiment that repeated administration of COX189 at 10 and 30mg/kg significantly inhibits the radiologically observed structural changes on day 20. COX189 at 30mg/kg is able to significantly inhibit the loss of bone mineral density compared to vehicle treated animals. Valdecoxib at 30mg/kg has no significant effect on either parameter.

CLAIMS

1. A method of treating cancer pain in a subject in need of such treatment which comprises administering to the subject an effective amount of a COX-2 inhibitor;
2. A method according to claim 1 in which the COX-2 inhibitor is a compound of formula I



wherein R is methyl or ethyl;

R₁ is chloro or fluoro;

R₂ is hydrogen or fluoro;

R₃ is hydrogen, fluoro, chloro, methyl, ethyl, methoxy, ethoxy or hydroxy;

R₄ is hydrogen or fluoro; and

R₅ is chloro, fluoro, trifluoromethyl or methyl;

a pharmaceutically acceptable salts thereof; or

a pharmaceutically acceptable prodrug esters thereof..

3. Use of a compound of formula I as defined in claim 2 (or pharmaceutically acceptable salt or prodrug ester thereof) for the preparation of a medicament, for use in the treatment of cancer pain.
4. Use of a compound of formula I as defined in claim 2 (or pharmaceutically acceptable salt or prodrug ester thereof) for the treatment of cancer pain.

5. A package comprising a compound of formula I as defined in claim 2 (or pharmaceutically acceptable salt or prodrug ester thereof) together with instructions for use in the treatment of cancer pain.
6. A method according to claim 2, or use according to claim 3, in which the compound of formula I is 5-methyl-2-(2-chloro-6-fluoroanilino)-phenylacetic acid or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable prodrug ester thereof.
7. A method according to claim 1 or use according to claim 3, for the treatment of bone cancer pain.
8. A method according to claim 1 or use according to claim 3, in which the compound of formula I is in the form of an oral composition or an injectable composition.
9. A method for the inhibition of bone loss, advantageously in cancer, which comprises administering an effective amount of a COX-2 inhibitor of formula I (or an ester or prodrug thereof) as defined in claim 2 to a subject in need of such treatment.
10. Use of a COX-2 inhibitor of formula I (or an ester or prodrug thereof) as defined in claim 2, for the preparation of a medicament for the inhibition of bone loss, in particular bone loss in cancer.

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
6 November 2003 (06.11.2003)

PCT

(10) International Publication Number
WO 2003/090737 A3

(51) International Patent Classification⁷: **A61K 31/196**,
C07C 229/42

WC1E 6BN (GB). **GONZALEZ, Isabel** [ES/GB]; 55 St
Barnabas Road, Cambridge, CB1 2BX (GB).

(21) International Application Number:
PCT/EP2003/004152

(74) Agent: **GRUBB, Philip**; Novartis AG, Corporate Intel-
lectual Property, CH-4002 Basel (CH).

(22) International Filing Date: 22 April 2003 (22.04.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0209257.5 23 April 2002 (23.04.2002) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LT, LU, LV, MA, MD, MK, MN, MX, NI, NO, NZ, OM,
PH, PL, PT, RO, RU, SC, SE, SG, SK, TJ, TM, TN, TR,
TT, UA, US, UZ, VC, VN, YU, ZA, ZW.

(84) Designated States (*regional*): Eurasian patent (AM, AZ,
BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE,
BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU,
IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR).

(71) Applicant (*for all designated States except AT, US*): **NO-
VARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel
(CH).

(71) Applicant (*for AT only*): **NOVARTIS PHARMA GMBH**
[AT/AT]; Brunner Strasse 59, A-1230 Viena (AT).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **FUJIMOTO**,
Roger, Aki [US/US]; 21 Molly Stark Drive, Morristown,
NJ 07960 (US). **URBAN, Laszlo** [GB/GB]; The Novartis
Institute for Medical Sciences, 5 Gower Place, London,


Published:

— with international search report

(88) Date of publication of the international search report:
1 April 2004

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: METHODS AND COMPOSITIONS FOR TREATMENT OF CANCER PAIN



(57) Abstract: The invention provides a method of treating cancer pain, e.g. bone cancer pain, in a subject in need of such treatment which comprises administering to the sub-
ject an effective amount of a COX-2 inhibitor; advantageously a compound of formula
(I); wherein R, R₁, R₂, R₃, R₄, and R₅ are as defined; a pharmaceutically acceptable salts
thereof; or a pharmaceutically acceptable prodrug ester thereof.

WO 2003/090737 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/04152

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/196 C07C229/42

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00 32189 A (GAO DANCHEN ;SEARLE & CO (US); MAZHARY AHMAD M (US); HLINAK ANTHON) 8 June 2000 (2000-06-08)	1,7,8
Y	claim 1 page 5, line 20 - line 21 page 9, line 30 - line 32 -----	1-8
X	WO 01 91750 A (HASSAN FRED ;BRUGGER ANDREW (US); FORBES JIM (US); GAO PING (US);) 6 December 2001 (2001-12-06)	1,7,8
Y	claim 1 * page 17, paragraph [0068] *	1-8
X	US 2002/013357 A1 (NADKARNI SREEKANT ET AL) 31 January 2002 (2002-01-31)	1,7,8
Y	claim 1 * page 1, paragraph [0004] * * page 3, paragraph [0049] * -----	1-8
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

28 July 2003

Date of mailing of the international search report

24.11.2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Beranová, P.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/04152

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 945 134 A (BOEHRINGER INGELHEIM PHARMA) 29 September 1999 (1999-09-29) claim 1 -----	1-8
Y	WO 99 11605 A (NOVARTIS AG) 11 March 1999 (1999-03-11) claims 1,8 -----	1-8
Y	MEDHURST S J ET AL: "A rat model of bone cancer pain." PAIN, vol. 96, no. 1-2, March 2002 (2002-03), pages 129-140, XP002249243 ISSN: 0304-3959 page 140, left-hand column, paragraph 3 -----	1,7,8

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 03/04152

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 9, 10
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.: --

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-8

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 9, 10

Present claims 1, 7 and 8 relate to a compound defined by reference to a desirable characteristic or property, namely "COX-2 inhibitor". The claims cover all compounds having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and disclosure within the meaning of Article 5 PCT for only a very limited number of such compounds. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the compound by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the compounds of formula (I) and rofecoxib, etoricoxib, celecoxib, valdecoxib and parecoxib as indicated in the description (page 1, 5th paragraph).

The applicant's attention is drawn to the fact that claims relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure. If the application proceeds into the regional phase before the EPO, the applicant is reminded that a search may be carried out during examination before the EPO (see EPO Guideline C-VI, 8.5), should the problems which led to the Article 17(2) declaration be overcome.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-8

The use of COX-2 inhibitors for the treatment of cancer pain

2. claims: 9, 10

The use of COX-2 inhibitors for the treatment of bone loss

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/04152

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0032189	A	08-06-2000	AT 225650 T	15-10-2002
			AU 748851 B2	13-06-2002
			AU 1838100 A	19-06-2000
			BG 104680 A	28-02-2001
			BR 9908030 A	28-11-2000
			CA 2319201 A1	08-06-2000
			CN 1288378 T	21-03-2001
			DE 69903404 D1	14-11-2002
			DE 69903404 T2	14-08-2003
			DK 1049467 T3	10-02-2003
			EA 3363 B1	24-04-2003
			EE 200000437 A	15-06-2001
			EP 1049467 A1	08-11-2000
			ES 2185412 T3	16-04-2003
			HR 20000434 A1	31-08-2000
			HU 0100867 A2	28-03-2002
			IL 136532 A	12-01-2003
			JP 2002531403 T	24-09-2002
			NO 20003815 A	29-09-2000
			NZ 505762 A	28-06-2002
			PL 341372 A1	09-04-2001
			PT 1049467 T	28-02-2003
			SI 1049467 T1	30-04-2003
			SK 11062000 A3	12-03-2001
			TR 200002207 T1	21-12-2000
			WO 0032189 A1	08-06-2000
			ZA 200002722 A	29-11-2000
WO 0191750	A	06-12-2001	AU 6501301 A	11-12-2001
			BR 0111186 A	10-06-2003
			CA 2408793 A1	06-12-2001
			CN 1444478 T	24-09-2003
			EP 1283710 A1	19-02-2003
			NO 20025631 A	22-01-2003
			WO 0191750 A1	06-12-2001
			US 2003134887 A1	17-07-2003
			US 2002028238 A1	07-03-2002
US 2002013357	A1	31-01-2002	AU 1805901 A	18-06-2001
			AU 1930301 A	18-06-2001
			AU 1931001 A	18-06-2001
			AU 1931101 A	18-06-2001
			AU 2041201 A	18-06-2001
			AU 750978 B2	01-08-2002
			AU 2057101 A	18-06-2001
			BG 105808 A	30-09-2002
			BG 105873 A	30-04-2002
			BR 0008058 A	26-03-2002
			BR 0008059 A	26-03-2002
			BR 0008060 A	05-02-2002
			BR 0008088 A	09-04-2002
			CA 2362673 A1	14-06-2001
			CA 2362675 A1	14-06-2001
			CA 2362815 A1	14-06-2001
			CA 2362816 A1	14-06-2001
			CN 1376146 T	23-10-2002
			CN 1379669 T	13-11-2002
			CN 1411447 T	16-04-2003

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/04152

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2002013357 A1		CZ 20012875 A3	13-02-2002
		CZ 20013162 A3	12-06-2002
		CZ 20013163 A3	12-06-2002
		CZ 20013210 A3	13-03-2002
		EA 3639 B1	28-08-2003
		EA 3612 B1	26-06-2003
		EA 3464 B1	26-06-2003
		EE 200100414 A	16-12-2002
		EE 200100419 A	16-12-2002
		EP 1175214 A2	30-01-2002
		EP 1165072 A2	02-01-2002
		EP 1150959 A1	07-11-2001
		EP 1150960 A1	07-11-2001
		HR 20010582 A1	31-08-2002
		HR 20010589 A1	31-08-2002
		HU 0200409 A2	29-06-2002
		HU 0200580 A2	28-11-2002
		HU 0201450 A2	28-12-2002
		JP 2003523954 T	12-08-2003
		JP 2003516353 T	13-05-2003
		JP 2003516394 T	13-05-2003
		NO 20013855 A	05-10-2001
		NO 20013858 A	08-10-2001
		NO 20013859 A	08-10-2001
		NO 20013868 A	03-10-2001
EP 0945134 A	29-09-1999	EP 0945134 A1	29-09-1999
		AU 762322 B2	26-06-2003
		AU 3417199 A	18-10-1999
		BR 9909192 A	05-12-2000
		CA 2326517 A1	07-10-1999
		CN 1295478 T	16-05-2001
		WO 9949867 A1	07-10-1999
		EP 1083902 A1	21-03-2001
		HU 0101246 A2	29-05-2002
		JP 2002509886 T	02-04-2002
		NO 20004823 A	22-11-2000
		NZ 507382 A	29-08-2003
		PL 342988 A1	16-07-2001
		TR 200002786 T2	22-01-2001
		ZA 200004178 A	20-02-2002
WO 9911605 A	11-03-1999	AT 237580 T	15-05-2003
		AU 743371 B2	24-01-2002
		AU 9534098 A	22-03-1999
		BR 9812010 A	12-12-2000
		CN 1268112 T	27-09-2000
		DE 69813570 D1	22-05-2003
		DK 1007505 T3	21-07-2003
		WO 9911605 A1	11-03-1999
		EP 1007505 A1	14-06-2000
		JP 2001514244 T	11-09-2001
		NO 20000943 A	25-02-2000
		NZ 502669 A	01-02-2002
		PL 338357 A1	23-10-2000
		RU 2186762 C2	10-08-2002
		SI 1007505 T1	31-10-2003
		SK 2472000 A3	12-09-2000

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/04152

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9911605	A	TR 200000447 T2	21-07-2000
		US 6291523 B1	18-09-2001
		US 2002183391 A1	05-12-2002
		US 6310099 B1	30-10-2001
		US 2002013369 A1	31-01-2002
		ZA 9807785 A	01-03-1999
		HU 0002514 A2	28-12-2000
